

On a vector bundle which cuts, bounces, embeds and measures waists

The vector bundles associated with the natural permutation representation $U_n := \mathbb{R}^n$, and the standard representation $W_n := \{(y_1, \dots, y_n) \in U_n : \sum y_i = 0\}$, of the symmetric group \mathfrak{S}_n over a free S -spaces X , $S \subseteq \mathfrak{S}_n$, are defined by

$$\xi = \xi_{(\mathfrak{S}_n, S, X)} : \quad U_n \longrightarrow X \times_S U_n \longrightarrow X/S,$$

$$\zeta = \zeta_{(\mathfrak{S}_n, S, X)} : \quad W_n \longrightarrow X \times_S W_n \longrightarrow X/S.$$

These bundles, in the case when $S = \mathfrak{S}_n$ and $X = F(M, n)$ is the classical configuration space of n pairwise distinct points on M , were originally studied and very efficiently used by Cohen, Cohen & Handel, Chisholm, Cohen, Mahowald & Milgram, Bödiger, Cohen & Taylor and many others. In the last decade, problems of

- the existence of convex measure partitions (the twisted Euler class of $\zeta_{(\mathfrak{S}_n, \mathfrak{S}_n, F(\mathbb{R}^d, n))}^{\oplus(d-1)}$),
- the existence of ℓ -skew embeddings (the dual Steifel–Whitney classes of $\xi_{(\mathfrak{S}_n, \mathfrak{S}_n, F(\mathbb{R}^d, n))}^{\oplus(d+1)}$),
- an estimation of waists of manifolds (the top Steifel–Whitney class of $\zeta_{(\mathfrak{S}_{2^m}, \mathfrak{S}_{2^m}^{(2)}, (S^{d-1})^{2^m-1})}^{\oplus(d-1)}$),
- counting periodic billiard trajectories (any non-zero characteristic class of $\zeta_{(\mathfrak{S}_p, \mathbb{Z}/p, G(\mathbb{R}^d, n))}$),

motivated Gromov, Ghomi & Tabachnikov, Karasev, Hubbard & Aronov, Crabb, and Blagojević, Lück & Ziegler, to start a new study on the properties of these vector bundles.

In this talk we go a bit deeper. Using an embedding of the product of spheres $(S^{d-1})^{2^m-1}$ into the configuration space $F(\mathbb{R}^d, 2^m)$, which is equivariant, only with respect to the action of a Sylow 2-subgroup \mathcal{S}_{2^m} of \mathfrak{S}_{2^m} , we first show that the cohomology ring $H^*(F(\mathbb{R}^d, 2^m)/\mathcal{S}_{2^m}; \mathbb{F}_2)$ of the unordered configuration space can be seen as a subring of the cohomology ring

$$H^*((S^{d-1})^{2^m-1}/\mathcal{S}_{2^m}; \mathbb{F}_2) \cong \mathbb{F}_2[V_{m,1}, \dots, V_{m,m}]/\langle V_{m,1}^d, \dots, V_{m,m}^d \rangle \oplus I^*(\mathbb{R}^d, 2^m),$$

where $I^*(\mathbb{R}^d, 2^m)$ is an ideal, and $\deg(V_{m,r}) = 2^r - 1$, $1 \leq r \leq m$.

In the next step we express the Stiefel–Whitney classes of the vector bundles $\xi := \xi_{(\mathfrak{S}_{2^m}, \mathcal{S}_{2^m}, F(\mathbb{R}^d, n))}$ in the language of $\mathrm{GL}_m(\mathbb{F}_2)$ -invariant Dickson polynomials. Then, using the subgroup $U_m(\mathbb{F}_2)$ of $\mathrm{GL}_m(\mathbb{F}_2)$, of upper triangular matrices with ones on the main diagonal, we realize the generators $V_{m,r}$ as $U_m(\mathbb{F}_2)$ invariants. Expressing recursively $\mathrm{GL}_m(\mathbb{F}_2)$ -invariants in terms of $U_m(\mathbb{F}_2)$ -invariants we explicitly identify the sequence $w_{2^m-2^0}(\xi), w_{2^m-2^1}(\xi), \dots, w_{2^m-2^{m-1}}(\xi)$ of Stiefel–Whitney classes in the polynomial part of the cohomology ring $H^*((S^{d-1})^{2^m-1}/\mathcal{S}_{2^m}; \mathbb{F}_2)$.

In this way, we made a step closer towards understanding the ideal generated by the Stiefel–Whitney classes of the vector bundle $\xi := \xi_{(\mathfrak{S}_n, \mathfrak{S}_n, F(\mathbb{R}^d, n))}$:

$$(w_1(\xi), w_2(\xi), \dots, w_{n-1}(\xi)) \in H^*(F(\mathbb{R}^d, n)/\mathfrak{S}_n; \mathbb{F}_2),$$

hoping to give complete answers to some of the questions listed above. This is a joint work with Frederick R. Cohen, Michael Crabb, Wolfgang Lück & Günter M. Ziegler

Одељење за математику је општи математички семинар намењен широкој публици. Предавања су прилагођена математичарима и онима који желе да то постану.

Зоран Петрић